

AD-A060 952

CALIFORNIA UNIV IRVINE
THEORY OF IMPURITY-SHIFTED INTERSUBBAND TRANSITIONS IN N-TYPE I--ETC(U)
1978 G M KRAMER, R F WALLIS
78-51


F/G 20/12
N00014-76-C-0121
NL

UNCLASSIFIED

| OF |
AD
A060 952



END
DATE
FILMED
2-79
DDC



DDC FILE COPY

AD A0 60 952

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER	2. DDT ACQUISITION NO.	3. RECIPIENT'S CATALOG NUMBER	12
4. TITLE (and Subtitle) Theory of Impurity-Shifted Intersubband Transitions in n-Type Inversion Layers in (100) Silicon.		5. TYPE OF REPORT & PERIOD COVERED FINAL rept.	R
6. AUTHOR(s) G. M./Kramer and R. F./Wallis		7. PERFORMING ORG. REPORT NUMBER 14 78-51	
8. PERFORMING ORGANIZATION NAME AND ADDRESS University of California, Irvine Irvine, CA 92717		9. CONTRACT OR GRANT NUMBER(s) 15 N00014-76-C-0121	
10. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research, Physics Program 800 N. Quincy St. Arlington, VA 22217		11. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 392-001	
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 125p.		13. REPORT DATE 11/1978	
14. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		15. NUMBER OF PAGES 4	
15. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		16. SECURITY CLASS. (of this report) Unclassified	
16. SUPPLEMENTARY NOTES		17. DECLASSIFICATION/DOWNGRADING SCHEDULE	
17. KEY WORDS (Continue on reverse side if necessary and identify by block number) Inversion layers, Silicon, Impurities, Optical Transition		DDC RECEIVED NOV 07 1978 E	
18. ABSTRACT (Continue on reverse side if necessary and identify by block number) Binding energies and variational wave functions are calculated as functions of electric field for electron impurity states split off from electric subbands of (100) silicon and silicon dioxide. The impurity-shifted intersubband transition energies are presented.			

DD FORM 1473 1 JAN 73 EDITION OF 1 NOV 65 IS OBSOLETE

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

387 30078 10 23 133

LB

SECRET

NOV 3 1973

ADDITION for	
WFO	White Section <input checked="" type="checkbox"/>
DOC	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	<input type="checkbox"/>
BY	
DISTRICT	SECURITY CODES
SIGNAL	

THEORY OF IMPURITY-SHIFTED INTERSUBBAND TRANSITIONS IN
N-TYPE INVERSION LAYERS ON (100) SILICON*

G. M. Kramer and R. F. Wallis

University of California, Irvine, California 92717, USA

Binding energies and variational wave functions are calculated as functions of electric field for electron impurity states split off from electric subbands of (100) silicon and silicon dioxide. The impurity-shifted intersubband transition energies are presented.

1. Introduction

The development of MOS devices has stimulated a great deal of interest in the basic properties of inversion layers at semiconductor boundaries. Of some importance are impurities and their associated electronic bound states. We have made a theoretical investigation of several bound states split off from the lowest three electric subbands of an n-type inversion layer on (100) silicon bounded by silicon dioxide.

2. Theoretical Development

We consider two contiguous, semi-infinite half-spaces, one of p-type silicon and the other of silicon dioxide, with a common boundary parallel to a (100) plane of the silicon. We assume that the potential energy of an electron undergoes an infinite discontinuous jump as the electron passes from the silicon into the silicon dioxide. In the region of the inversion layer, there is an electric field which, to a good first approximation¹, can be taken to be constant.

We assume that an impurity ion of charge $+Ze$ is located at the boundary between the silicon and the oxide. For this situation², the lowest lying bound states associated with a (100) interface on silicon are derived from the constant energy ellipsoids whose major axes are perpendicular to the interface.

The Hamiltonian is taken to be

$$H = H_0 + H_1 \quad (1)$$

where

$$H_0 = -\gamma \frac{\partial^2}{\partial z^2} + \frac{\delta}{z} + \kappa \epsilon z, \quad z \geq 0, \quad (2)$$

$$H_1 = -\frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial y^2} + U(\vec{r}), \quad (3)$$

$\gamma = m_t/m_l$, $\delta = (\epsilon_2 - \epsilon_1)/4Z\epsilon_2$, $\kappa = \hbar^4(\epsilon_1 + \epsilon_2)^3/4Z^3 m_t^2 e^5$, ϵ is the electric field in esu, m_t and m_l are the transverse and longitudinal effective masses, and ϵ_1 and ϵ_2 are the dielectric constants of SiO_2 and silicon, respectively. In Eq. (1), energy is measured in units of the effective Rydberg, $Ry^* = 2m_t e^4 Z^2 / \hbar^2 (\epsilon_1 + \epsilon_2)^2$, and length in units of the effective Bohr radius, $a_0^* = \hbar^2 (\epsilon_1 + \epsilon_2) / 2Z m_t e^2$. For the silicon- SiO_2 system, $Ry^* = 42.3$ meV and $a_0^* = 21.8$ Å. The impurity ion is taken to be at the silicon- SiO_2 interface and screening is neglected, so $U(\vec{r})$ has the Coulomb form

$$U(\vec{r}) = -\frac{2}{|\vec{r}|}. \quad (4)$$

The term involving δ in Eq. (2) represents the interaction of the electron with its own image.

In the limit of high electric fields, an adiabatic theorem applies and the impurity wave function has the form

$$\psi(\vec{r}) = \chi_m^{(n)}(x, y) f_n(z) \quad (5)$$

where $f_n(z)$ is an eigenfunction of H_0 given by Eq. (2) and corresponds to the electric subband of interest. We consider three subbands and choose the $f_n(z)$ to be variational functions of the Fang-Howard³ type

$$f_0(z) = A_0 z \exp(-\frac{1}{2} b_0 z) \quad (6a)$$

$$f_1(z) = A_1 z (1 - \alpha_1 z) \exp(-\frac{1}{2} b_1 z) \quad (6b)$$

$$f_2(z) = A_2 z (1 + \alpha_2 z + \beta_2 z^2) \exp(-\frac{1}{2} b_2 z) \quad (6c)$$

The quantities b_n are variational parameters, the A_n are normalization constants, and α_1 , α_2 , and β_2 are chosen so that the functions $f_n(z)$ are orthogonal on the interval $0 \leq z \leq \infty$. We take $\chi_m^{(n)}(x, y)$ to have either of two forms

$$\chi_0^{(n)}(x, y) = B_0 \exp[-\frac{1}{2} a_0^{(n)} (x^2 + y^2)^{\frac{1}{2}}] \quad (7a)$$

$$\chi_{\pm 1}^{(n)}(x, y) = B_1(x \pm iy) \exp[-\frac{1}{2} a_1^{(n)}(x^2 + y^2)^{\frac{1}{2}}] \quad (7b)$$

where the B_1 are normalization constants and the $a_1^{(n)}$ are variational parameters. The parameters $a_1^{(n)}$ and b_n are varied independently to minimize the expectation value of the Hamiltonian, $\langle H \rangle$. The expectation value of H_0 , $\langle H_0 \rangle$, is also minimized separately with respect to the b_n to give the continuum energy. The impurity binding energy, E_B , is then given by

$$E_B = \langle H_0 \rangle - \langle H \rangle \quad (8)$$

3. Results and Discussion

The impurity binding energies have been calculated for electric fields between 10^2 and 10^9 esu for the three lowest subbands and $m=0, \pm 1$. The results are plotted in Fig. 1.




Fig. 1. Impurity state binding energies versus field

The impurity binding energy decreases as one goes from the lowest subband to successively higher subbands for both the $m=0$ and $m=\pm 1$ bound states. For a given bound state associated with a given subband, the binding energy increases with increasing electric field and approaches the two-dimensional value in the limit of infinite field.

Electric dipole transitions between subbands are allowed transitions.³ In Fig. 2 are given our calculated impurity shifted transition energies for the 0-1 and 0-2 inter-

subband transitions when $m=0$. The transition energies are shifted to higher energies when the electron becomes localized at the impurity. At a field of 10^2 esu, the impurity shifted 0-1 transition is 70 percent higher than the unshifted transition and is only 15 percent lower than the 0-2 unshifted transition. The latter result is in qualitative agreement with recent experimental data.⁴ As the electric field increases, the impurity shifted 0-1 transition drops back toward the 0-1 free electron transition and away from the 0-2 free electron transition.




Fig. 2. Intersubband transition energies for bound ($m=0$) and free states

References

- * Work supported in part by the Office of Naval Research
- 1. Stern F and Howard W E 1967 Phys. Rev. **163** 816
- 2. Karpushin A A 1970 Soviet Physics-Solid State **11** 1748
- 3. Kneschaurek P and Koch J F 1977 Phys Rev **B16** 1590
- 4. McCombe B D and Shafer D E to be presented at this Conference and private communication.

ACCESSION BY	
DATE	WHA Section <input checked="" type="checkbox"/>
ORG	JOIT Section <input type="checkbox"/>
REPRODUCED	<input type="checkbox"/>
NOT REPRODUCED	
BY	
DISTRIBUTION AVAILABILITY CODES	
DECL	AVAIL. NOT RE. SPECIAL
A	